

6 | Momentum

IB Physics Content Guide

Big Ideas

- The total momentum of an isolated system is always constant
- The force on an object when speeding up or slowing down can be affected by changing the time for the force
- The impulse of a collision is equal to the change in momentum

Content Objectives

6.1 – Conservation of Momentum

 p. 92, 96-101

I can define and calculate momentum			
I can calculate “before” and “after” momentums for multiple objects			
I can use the conservation of momentum to solve for missing variables in linear collisions			
I can describe the process required for explosion, hit and bounce, and hit and stick scenarios			
I can describe the difference between elastic and non-elastic collisions			
I can describe how energy is not always conserved within a system			
I can calculate the amount of energy retained in a non-elastic collision			

6.2 – Momentum and Impulse

 p. 93-95

I can describe the meaning of impulse and how it is related to momentum change			
I can use impulse and momentum to solve for an unknown in a collision problem			
I can conceptually describe how to decrease the force experienced in a collision			
I can determine the impulse of a collision from a force vs time graph			

6 | Momentum

Shelving Guide

	Variable Symbol	Unit
Momentum		
Mass		
Velocity		
Time		
Kinetic Energy		
Impulse	Impulse	

Data Booklet Equations:

$$p = mv$$

$$F = \frac{\Delta p}{\Delta t}$$

$$E_K = \frac{p^2}{2m}$$



$$\text{Impulse} = F\Delta t = \Delta p$$

Conservation of Energy Problems

Types of Collisions

Elastic	
Inelastic	

Calculating Impulse

<i>Constant force:</i>	 <p>A diagram showing a silver car on a grey surface. A blue arrow points to the right from the front of the car, labeled "5000 N". To the right of the car, a stopwatch icon is shown with the text "8.9 s" next to it. Two vertical blue lines mark the start and end of the time interval.</p>
<i>Varying Force:</i>	 <p>A graph with Force (N) on the vertical axis and Time (s) on the horizontal axis. The vertical axis ranges from 0 to 13 with major grid lines every 1 unit. The horizontal axis ranges from 0 to 10 with major grid lines every 1 unit. A red line starts at (0,0), rises linearly to a peak at (4,8), and then falls linearly to (10,0).</p>

Impulse-Momentum Equation

Collision Safety

Explain (using impulse, force, and time) how to decrease the force acting on an object undergoing a collision: